

LOWER BOUNDS FOR THE CANONICAL HEIGHT ON ELLIPTIC CURVES OVER ABELIAN EXTENSIONS

MATTHEW H. BAKER

ABSTRACT. Let K be a number field and let E/K be an elliptic curve. If E has complex multiplication, we show that there is a positive lower bound for the canonical height of non-torsion points on E defined over the maximal abelian extension K^{ab} of K . This is analogous to results of Amoroso-Dvornicich and Amoroso-Zannier for the multiplicative group. We also show that if E has non-integral j -invariant (so that in particular E does not have complex multiplication), then there exists $C > 0$ such that there are only finitely many points $P \in E(K^{\text{ab}})$ of canonical height less than C . This strengthens a result of Hindry and Silverman.

1. INTRODUCTION

Let K be a number field, and let K^{ab} denote the maximal abelian extension of K . Let E/K be an elliptic curve, and let $\hat{h} : E(\bar{K}) \rightarrow \mathbb{R}$ be the usual canonical height function on $E(\bar{K})$.

Suppose that E has complex multiplication by an order in the imaginary quadratic field k . If K is any number field containing k , the theory of complex multiplication tells us that all of the torsion points of $E(\bar{K})$ (which are exactly the points with canonical height zero) are defined over the maximal abelian extension K^{ab} of K . By analogy with recent results of Amoroso–Dvornicich and Amoroso–Zannier for the multiplicative group (see Theorem 1.3 below), one might ask whether or not the canonical height function, when restricted to $E(K^{\text{ab}})$, takes arbitrarily small positive real values. The following theorem answers this question.

Theorem 1.1. *Let K be a number field, and let E/K be an elliptic curve having complex multiplication. Then there exists an effectively computable real constant $C > 0$ (depending on E/K) such that $\hat{h}(P) \geq C$ for all non-torsion points $P \in E(K^{\text{ab}})$.*

Remark 1.2. We will prove a somewhat more precise statement (Theorem 4.1) in Section 4.

Theorem 1.1 is analogous to the following result for heights on the multiplicative group $\mathbf{G}_m$, which (although not stated in precisely this form) is proved in [1] and [2]:

Theorem 1.3. *Let h denote the usual absolute logarithmic Weil height on $\overline{\mathbb{Q}}$. Let K be a number field, and let K^{ab} be the maximal abelian extension of K . Then there exists an effectively computable constant $C > 0$ (depending only on K) such that if $\alpha \in K^{\text{ab}}$ is not equal to zero or a root of unity, then $h(\alpha) \geq C$.*

Date: Revised on December 16, 2002.

The author's research was supported in part by an NSF Postdoctoral Research Fellowship.

Now suppose that E does not have complex multiplication. As a complement to Theorem 1.1, we prove the following result:

Theorem 1.4. *Suppose K is a number field and E is an elliptic curve with non-integral j -invariant. Then there exists a constant $C > 0$ such that $\hat{h}(P) \geq C$ for all but finitely many points $P \in E(K^{\text{ab}})$.*

Remark 1.5. The hypothesis that j_E is non-integral implies that E does not have complex multiplication. The elliptic curves not covered by either Theorem 1.1 or Theorem 1.4 are exactly those which have everywhere potential good reduction but do not have complex multiplication.

Remark 1.6. Theorem 1.4 generalizes a result of Hindry-Silverman (Corollary 0.3 of [6]), which says that under the hypotheses of Theorem 1.4, there exist effectively computable constants $C, D > 0$ such that any point $P \in E(K^{\text{ab}})$ with $\deg(P) \geq D$ satisfies the inequality

$$\hat{h}(P) \geq \frac{C}{\deg(P)^{2/3}}.$$

Silverman also proved in [17] that if E/K is an elliptic curve without complex multiplication (over $\bar{K}$), then there exists an effectively computable constant $C > 0$ such that every non-torsion point $P \in E(K^{\text{ab}})$ satisfies

$$\hat{h}(P) \geq \frac{C}{\deg(P)^2}.$$

Remark 1.7. The proof of Theorem 1.4 does not provide an effective algorithm for computing the constant C in the statement of the theorem.

We conclude this section by placing Theorems 1.1 and 1.4 in a more general context.

Let G be a variety defined over a field K , and let L/K be a field extension. Suppose that $\hat{h} : G(L) \rightarrow \mathbb{R}$ is a function taking nonnegative values and that $S \subseteq G(L)$. We say that the pair $(S, \hat{h})$ (or the set S , when the function $\hat{h}$ is understood) has the *discreteness property* if there exists a constant $C > 0$ such that $\hat{h}(P) \geq C$ whenever $P \in S$ and $\hat{h}(P) > 0$. We say that the pair $(S, \hat{h})$ has the *strong discreteness property* if in fact we have

$$\liminf_{P \in S} \hat{h}(P) > 0,$$

i.e., if there exists a constant $C > 0$ such that $\hat{h}(P) \geq C$ for all but finitely many $P \in S$.

An example to keep in mind is the case where $G = A$ is an abelian variety defined over a number field K and $\hat{h} : A(\bar{K}) \rightarrow \mathbb{R}$ is the canonical height function corresponding to some symmetric ample line bundle on A . It is well known (see [11, Section 4], for example) that whether or not $(S, \hat{h})$ has the discreteness property (resp. the strong discreteness property) is independent of which canonical height function one chooses. Another example is where $G = \mathbf{G}_m^n$ and $\hat{h} : G(\bar{\mathbb{Q}}) \rightarrow \mathbb{R}$ is the function given $\hat{h}(x_1, \dots, x_n) = h(x_1) + \dots + h(x_n)$, where $h : \bar{\mathbb{Q}} \rightarrow \mathbb{R}$ is the usual logarithmic Weil height.

Theorem 1.3 implies that $\mathbf{G}_m^n(K^{\text{ab}})$ has the discreteness property for all number fields K . Theorem 1.1 can be rephrased as saying that if E/K is an elliptic curve

with complex multiplication, then $E(K^{\text{ab}})$ has the discreteness property. Similarly, Theorem 1.4 says that if E/K is an elliptic curve with non-integral j -invariant, then $E(K^{\text{ab}})$ has the strong discreteness property.

We mention some known results related to Theorems 1.1 and 1.4. Concerning torsion points of abelian varieties defined over abelian extensions, J. Zarhin [23] (see also Ruppert [13]) has proved the following, using deep finiteness results of Faltings ([5]):

Theorem 1.8. *Let A be an abelian variety defined over a number field K . Then the torsion subgroup $A_{\text{tor}}(K^{\text{ab}})$ of $A(K^{\text{ab}})$ is finite iff A has no abelian subvariety with complex multiplication over K .*

There is also the following result due to S. Zhang, which is a consequence (using Weil restriction of scalars) of the equidistribution theorems proved in [24] and [20]:

Theorem 1.9 (Zhang). *Let K be a number field, and let A/K be an abelian variety. If $L \supseteq K$ and L is a finite extension of a totally real field, then $A(L)$ has the strong discreteness property.*

Since the maximal cyclotomic extension $K^{\text{cycl}} := K(\mu_\infty)$ is a finite extension of the totally real field $\mathbb{Q}(\mu_\infty)^+$, Zhang's result implies in particular:

Corollary 1.10. *Let A/K be an abelian variety. Then $A(K^{\text{cycl}})$ has the strong discreteness property.*

The weaker fact that the torsion subgroup $A_{\text{tor}}(K^{\text{cycl}})$ is finite is originally due to Ribet [12].

In light of the above-mentioned results, we propose the following conjecture.

Conjecture 1.11. *Let K be a number field, and let A/K be an abelian variety. Then:*

1. *$A(K^{\text{ab}})$ has the discreteness property.*
2. *If A does not contain an abelian subvariety having complex multiplication over K , then $A(K^{\text{ab}})$ has the strong discreteness property.*

Note that Theorems 1.1 and 1.4 are special cases of part (1) of this conjecture. Together with Corollary 1.10, these results imply that Conjecture 1.11 is true when $A = E$ is an elliptic curve and any one of the following three hypotheses is satisfied:

- E has complex multiplication
- j_E is non-integral
- $K = \mathbb{Q}$.

We note that if $P_0 \in A(K)$ is a nontorsion point and $P_n \in A(\bar{K})$ satisfies $2^n P_n = P_0$, then $\hat{h}(P_n) = \frac{1}{4^n} \hat{h}(P_0) \rightarrow 0$ as $n \rightarrow \infty$. This is the simplest way to construct a sequence of nontorsion points in $A(\bar{K})$ whose heights tend toward zero. Conjecture 1.11 is therefore related to the principle that when n is large, the points P_n tend to be defined over highly non-abelian extensions of K .

To conclude this introduction, we pose the following question (which one could also ask in the context of semiabelian varieties):

Question: If A/K is an abelian variety and L is the extension of K generated by all torsion points of $A(\bar{K})$, is it true that $A(L)$ has the discreteness property?

2. BACKGROUND FACTS ON CANONICAL HEIGHTS

If K is a number field, we let M_K denote the set of places (i.e., equivalence classes of absolute values) of K . We will denote the archimedean places by M_K^∞ and the non-archimedean ones by M_K^0 . If $v \in M_K$, we let $|\cdot|_v$ be the unique absolute value in the equivalence class v which extends the usual absolute value on $\mathbb{Q}_v$. We also let $n_v = [K_v : \mathbb{Q}_v]$ denote the local degree at v , and we let $d_v = \frac{[K_v : \mathbb{Q}_v]}{[K : \mathbb{Q}]}$ denote the local degree divided by the global degree.

We have:

Lemma 2.1. *If L is a finite extension of the number field K and v is a fixed place of K , then*

$$\sum_{w \in M_L, w|v} d_w = d_v.$$

PROOF. Using the well-known fact that $\sum_{w|v} [L_w : K_v] = [L : K]$, we have

$$\sum_{w|v} \frac{[L_w : \mathbb{Q}_w]}{[L : \mathbb{Q}]} = \sum_{w|v} \frac{[L_w : K_v][K_v : \mathbb{Q}_v]}{[L : \mathbb{Q}]} = \frac{[K_v : \mathbb{Q}_v]}{[K : \mathbb{Q}]}$$

as desired. ■

Let E/K be an elliptic curve. It is well known (see [19, Chapter VI] for definitions and further properties) that the canonical height function $\hat{h} : E(\bar{K}) \rightarrow \mathbb{R}$ has the following properties:

- $\hat{h}(nP) = n^2 \hat{h}(P)$ for all $P \in E(\bar{K})$ and all integers n .
- $\hat{h}(P) \geq 0$ for all $P \in E(\bar{K})$, and $\hat{h}(P) = 0$ if and only if P is a torsion point.
- (Northcott's theorem) Given any constant $M > 0$ and a natural number d , the set of points $P \in E(\bar{K})$ such that $\hat{h}(P) \leq M$ and $[K(P) : K] \leq d$ is finite.
- (Parallelogram law) For all $P, Q \in E(\bar{K})$,

$$\hat{h}(P + Q) + \hat{h}(P - Q) = 2\hat{h}(P) + 2\hat{h}(Q).$$

The canonical height $\hat{h}$ has a decomposition into Néron local height functions λ_v , one for each $v \in M_K$. More precisely, for all $P \in E(K), P \neq 0$ we have

$$\hat{h}(P) = \sum_{v \in M_K} n_v \lambda_v(P),$$

where $\lambda_v : E(K_v) - \{0\} \rightarrow \mathbb{R}$ is defined and normalized as in [19, Chapter VI]. Note that in general, the Néron local height functions can take negative values, although if v is a place of good reduction for E , then $\lambda_v(P)$ is always nonnegative. In fact, suppose E has good reduction at v and that $\mathcal{E}$ is a minimal Weierstrass model over the ring of integers R_v in the completion K_v . Then if v corresponds to the prime ideal $\mathfrak{p} = (\pi)$ of R_v and $P \in E(K_v) - \{0\}$, we have

$$\lambda_v(P) = \log |\pi|_v^{m(P)},$$

where $m(P)$ is the largest integer $m \geq 0$ such that $P \equiv 0 \pmod{\pi^m}$.

The Néron local heights behave functorially with respect to finite extensions of K : if L/K is a finite extension and $P \in E(K)$, then

$$l_v(P) = \sum_{w|v} \frac{[L_w : K_v]}{[L : K]} l_w(P).$$

In particular, for each $v \in M_K$, λ_v extends to a Néron local height function $l_v : E(\overline{K_v}) - \{0\} \rightarrow \mathbb{R}$.

In order to prove Theorems 1.1 and Theorem 1.4, we will need known lower bounds for certain average values of Néron local height functions. We consider the archimedean case first.

Lemma 2.2. *Let $E/\mathbb{C}$ be an elliptic curve with j -invariant j_E , and let $l : E(\mathbb{C}) - \{0\} \rightarrow \mathbb{R}$ denote the associated Néron local height function. Let $P_1, \dots, P_N$ be distinct points of $E(\mathbb{C})$. Then*

$$\liminf_{N \rightarrow \infty} \frac{1}{N(N-1)} \sum_{i \neq j} l(P_i - P_j) \geq 0.$$

More explicitly, we have

$$\sum_{i \neq j} l(P_i - P_j) \geq -\frac{1}{12} N \log N - \frac{1}{12} N \log^+ |j_E| - \frac{11}{3}.$$

PROOF. This follows from a result of Elkies which is proved in a slightly different form in [8, Theorem VI.5.1]. The necessary modifications to get the form quoted here are discussed in [7, Lemma 3]. $\blacksquare$

It is clear from the definition of l that there exists a real constant C_1 such that $l(P) \geq -C_1$ for all $P \in E(\mathbb{C}) - \{0\}$. Taking $N = 2$ in the above lemma, we obtain the following explicit admissible value for C_1 :

Corollary 2.3. *Let $E/\mathbb{C}$, l be as above. Then if $P \in E(\mathbb{C}) - \{0\}$, we have*

$$l(P) \geq -C_1,$$

where $C_1 = \frac{1}{6}(C_2 + \log^+ |j_E|)$ and C_2 is a universal constant which can be taken to be $\log 2 + \frac{22}{3}$.

We now turn to the special case where v is a non-archimedean place of K at which E has split multiplicative reduction, so that in particular the j -invariant of E is non-integral at v . Let $\nu = \text{ord}_v(j_E^{-1}) > 0$, and note also for future reference that $\log |j_E|_v > 0$.

Let $\mathcal{E}$ be the Néron model of E over the ring of integers of K_v , and let $\tilde{E}$ be its special fiber. We denote by Φ the group of connected components of $\tilde{E}$ and we fix an isomorphism

$$\Phi \xrightarrow{\sim} \frac{(1/\nu)\mathbb{Z}}{\mathbb{Z}}$$

(see [19, Corollary IV.9.2]).

Let $l_v : E(K_v) - \{0\} \rightarrow \mathbb{R}$ be the Néron local height function at v . Tate's explicit formula for l_v implies that if r_v denotes the natural reduction homomorphism

$$r_v : E(K_v) \rightarrow \Phi \xrightarrow{\sim} \frac{(1/\nu)\mathbb{Z}}{\mathbb{Z}},$$

then

$$l_v(P) \geq \frac{1}{2} \mathbf{B}_2(r_v(P)) \log |j_E|_v$$

for all nonzero points $P \in E(K_v)$. Here $\mathbf{B}_2(t)$ is the periodic second Bernoulli polynomial $\{t\}^2 - \{t\} + \frac{1}{6}$. Note in particular that if $P \in E(K_v) - \{0\}$ reduces to the identity component of $\tilde{E}$ then

$$l_v(P) \geq \frac{1}{12} \log |j_E|_v > 0.$$

Using this description and a Fourier averaging technique, one can prove the following analogue of Lemma 2.2 (see [6, Proposition 1.2]):

Lemma 2.4. *Let v be a finite place of K such that j_E is non-integral at v , and let $P_1, \dots, P_N \in E(K_v)$ be distinct points. Let $\nu = \text{ord}_v(j_E^{-1})$. Then*

$$\liminf_{N \rightarrow \infty} \frac{1}{N(N-1)} \sum_{i \neq j} l_v(P_i - P_j) \geq 0.$$

More explicitly, we have

$$\sum_{i \neq j} \lambda_v(P_i - P_j) \geq \frac{1}{12} \log |j_E|_v \left(\left(\frac{N}{\nu} \right)^2 - N \right).$$

3. PRELIMINARY LEMMAS

In this section, we prove some lemmas which will be needed for the proof of Theorem 1.1. The first lemma is well-known, and concerns the formal group of an abelian variety with ordinary reduction. For lack of a suitable reference, we sketch a proof.

Lemma 3.1. *Let R be a strictly henselian discrete valuation ring with fraction field K of characteristic zero and residue field k of characteristic $p > 0$. Let A/K be an abelian variety with good ordinary reduction, and let $\mathcal{A}/R$ be its Néron model. Let $\hat{\mathcal{A}}$ be the formal group (i.e., the formal completion along the identity section) of $\mathcal{A}$. Then $\hat{\mathcal{A}}$ is isomorphic over R to $(\hat{\mathbf{G}}_m)^g$, where $g = \dim A$ and $\hat{\mathbf{G}}_m$ is the formal group of $\mathbf{G}_m$.*

PROOF. (Sketch) Let G/R be the p -divisible group of $\mathcal{A}$, so that by the Serre-Tate correspondence, the formal group $\hat{\mathcal{A}}$ is the connected component G^0 of G . Since A has good ordinary reduction and k is separably closed, G contains a constant étale p -divisible group isomorphic to $(\mathbb{Q}_p/\mathbb{Z}_p)^g$.

On the other hand, let $G^\vee$ be the p -divisible group of the dual abelian variety $A^\vee$. Then G and $G^\vee$ are canonically Cartier dual to one another (see [10]). Additionally, A and $A^\vee$ are isogenous over $\bar{K}$, so that $A^\vee$ also has good ordinary reduction. Therefore $G^\vee$ contains a copy of $(\mathbb{Q}_p/\mathbb{Z}_p)^g$, and by Cartier duality, it follows that G contains a split multiplicative p -divisible group isomorphic to $(\mathbf{G}_m)^g$.

It follows by dimension considerations that $G \cong (\mathbb{Q}_p/\mathbb{Z}_p)^g \times (\mathbf{G}_m)^g$ and therefore $\hat{\mathcal{A}} \cong (\hat{\mathbf{G}}_m)^g$ as desired. ■

Our proof of Theorem 1.1 will follow the same basic outline as the proof of [1, Theorem 1], replacing the map $x \mapsto x^p$ on $\mathbf{G}_m$ for a suitable prime number p by a

canonical lift of Frobenius on E . So we now recall some facts concerning canonical lifts.

As in the introduction, let K be a number field with ring of integers R , and let E/K be an elliptic curve with complex multiplication by an order R' in the imaginary quadratic field k . Let v be a non-archimedean place of K , corresponding to a prime ideal $\mathfrak{p}$ of R of residue characteristic $p > 2$, and let K_v (resp. R_v) denote the v -adic completion of K (resp. R). Throughout this section, we make the additional assumptions that E has good ordinary reduction at v and that p does not divide the conductor of R' . Our hypotheses on p ensure that there exists a canonical lift $F \in \text{End}_{\bar{K}}(E)$ of the Frobenius endomorphism of $\tilde{E}$ (see e.g. Chapter 13, Theorems 12–14 of [9]). The endomorphism F is an isogeny of degree p , and by replacing K by a finite extension if necessary, we can and will assume that F is defined over K . Also, the universal property of the Néron model (see [3, 1.1]) ensures that F extends to an endomorphism $F : \mathcal{E} \rightarrow \mathcal{E}$.

Let $\mathcal{E} : y^2 = f(x)$ be a minimal Weierstrass model for E/R_v , and let $\tilde{E}$ be the special fiber of $\mathcal{E}$. We denote by $\hat{\mathcal{E}}$ the formal group of $\mathcal{E}$.

Concerning the canonical lifting of Frobenius, we have the following lemma:

Lemma 3.2. *With notation and hypotheses as above (so that, in particular, E/K is an elliptic curve with complex multiplication and good ordinary reduction at v), let I be the inertia subgroup of $\text{Gal}(\bar{K}_v/K_v)$. Then $E[F^n] \cong \mu_{p^n}$ as I -modules for all integers $n \geq 1$.*

PROOF. We claim that $\mathcal{E}[F^n] = \mathcal{E}[p^n]^0$ as finite flat group schemes over R for all $n \geq 1$. One way to see this is as follows. By a theorem of Deligne (see [21, pp. 144–145]), $\mathcal{E}[F^n]$ is killed by its order p^n . But since $\mathcal{E}[F^n]$ is connected, it is a closed subscheme of $\mathcal{E}[p^n]^0$. Since E is ordinary, a consideration of degrees forces the equality $\mathcal{E}[F^n] = \mathcal{E}[p^n]^0$, proving the claim.

By the Serre-Tate correspondence, the formal group of $\mathcal{E}$ is the connected component of the p -divisible group of $\mathcal{E}$. It follows that $\mathcal{E}[F^n] = \mathcal{E}[p^n]^0 = \hat{\mathcal{E}}[p^n]$, and the result now follows from Lemma 3.1. $\blacksquare$

We also note for future reference the fact that for all $P \in E(\bar{K})$, $\hat{h}(FP) = p\hat{h}(P)$. This follows from the more general fact that for all nonzero $\psi \in \text{End}(E)$, we have $\hat{h}(\psi(P)) = \deg(\psi)\hat{h}(P)$ for all $P \in E(\bar{K})$ (see e.g. [15, Exercise 3.6.3]).

As a first step toward proving Theorem 1.1, we consider the special case where L/K is an abelian extension which is *unramified* above v . In this case, let $\sigma \in \text{Gal}(L/K)$ be the Frobenius element at v . (This is well-defined since L/K is abelian.)

Lemma 3.3. *Let L/K be an abelian extension, unramified above v , and suppose $P \in E(L)$ is not a torsion point. Then $FP \neq \sigma P$, and*

$$l_w(FP - \sigma P) \geq \log p$$

for all places w of L dividing v .

PROOF. Suppose $FP = \sigma P$. Then by induction, we have $F^k P = \sigma^k P$ for all $k \geq 1$. If M denotes the order of σ in $\text{Gal}(L/K)$, then we have $(F^M - 1)P = 0$. But $F^M - 1$ is an isogeny, which contradicts the fact that P is not a torsion point.

For the second part of the lemma, we note that for $Q \in E(L)$, $Q \neq 0$, we have $l_w(Q) > 0$ if and only if Q is in the kernel of reduction mod w . Also, since L_w/K_v

is unramified, the inequality $l_w(Q) > 0$ implies that $l_w(Q) \geq \log p$. The lemma follows from this last observation, since $FP - \sigma P$ is in the kernel of reduction mod w by the definition of F and σ . $\blacksquare$

Remark 3.4. Conversely, if $P \in E(L)$ is a torsion point then $FP = \sigma P$. So Lemma 3.3 gives a useful criterion for distinguishing non-torsion points from torsion points.

We now consider the special case of Theorem 1.1 in which $K_v = \mathbb{Q}_p$ and $L_w = \mathbb{Q}_p(\zeta_m)$ for some positive integer m divisible by p . Let $\tau \in \text{Gal}(L/K)$ be a generator of $\text{Gal}(\mathbb{Q}_p(\zeta_m)/\mathbb{Q}_p(\zeta_{m/p}))$, which can be identified with a subgroup of the inertia group I of $\text{Gal}(L_w/K_v)$.

The following elementary observation is similar to [1, Lemma 2(1)]:

Lemma 3.5. *With τ, m as above, we have $\tau(\alpha)^p \equiv \alpha^p \pmod{p}$ for all $\alpha \in \mathbb{Z}_p[\zeta_m]$.*

PROOF. Write $\alpha = h(\zeta_m)$ for some polynomial $h \in \mathbb{Z}_p[x]$. Then

$$\alpha^p \equiv h(\zeta_m^p) \equiv h(\tau \zeta_m^p) \equiv \tau \alpha^p \pmod{p}.$$

$\blacksquare$

The next lemma will allow us to prove Theorem 1.1 via induction on the ramification of L/K . It is modeled after Lemma 2(2) of [1].

Lemma 3.6. *Let $m = p^k m'$ be an integer with $(m', p) = 1$, let L/K be a finite abelian extension, and let w be a place of L lying over the place v of K . Assume that $L_w \cong \mathbb{Q}_p(\zeta_m)$, $K_v \cong \mathbb{Q}_p$, and that τ is a generator of $\text{Gal}(\mathbb{Q}_p(\zeta_m)/\mathbb{Q}_p(\zeta_{m/p}))$. Assume furthermore that $L \supseteq K(E[F^k])$. Suppose $P \in E(L)$ is a non-torsion point, and let $P' = F\tau P - FP$.*

1. *If $P' \neq 0$, then $l_w(P') \geq \log p$.*
2. *If $P' = 0$, then there exists a torsion point $T \in E[F^k]$ such that $P + T$ is fixed by τ .*

PROOF. For the first statement, we view P as a point of $E(\mathbb{Q}_p(\zeta_m))$. Let $\mathbb{Q}_p^{\text{nr}}$ denote the completion of the maximal unramified extension of $\mathbb{Q}_p$, and let I be the inertia group of $L_w/\mathbb{Q}_p$. According to Lemma 2 of [16], the reduction map gives an isomorphism

$$E[m](\mathbb{Q}_p^{\text{nr}}) \cong \tilde{E}[m](\overline{\mathbb{F}}_p).$$

It follows that there exists a torsion point $S \in E[m](\mathbb{Q}_p^{\text{nr}})$ such that $P - S$ is not in the kernel of reduction. Since τ fixes torsion points unramified at v and F is defined over K , we have

$$(\tau F - F)P = (\tau F - F)(P - S),$$

and therefore we may assume without loss of generality for the first part of the lemma that P is not in the kernel of reduction. Therefore we may write $P = (x, y)$ with $x, y \in \mathbb{Z}_p[\zeta_m]$.

Since τ is in the inertia group I , it follows that FP and τFP are congruent modulo the maximal ideal $\mathcal{M}$ of $\mathbb{Z}_p[\zeta_m]$. We claim that they are in fact congruent mod $\mathcal{M}^e = (p)$, which will establish part (1) of the lemma.

To see this, simply note that in terms of Weierstrass coordinates, we have $\tau FP \equiv (\tau x^p, \tau y^p)$ and $FP \equiv (x^p, y^p) \pmod{p}$. The claim therefore follows from Lemma 3.5.

For the second part of the lemma, suppose that $F\tau P = FP$. Then $\tau P - P \in E[F]$, which is a cyclic group of order p , so we have $\tau P - P = T'$, where T' is a

torsion point of order dividing p . If $T' = 0$ then we can take $T = 0$ in the statement of the lemma, so without loss of generality, we can suppose that T' has order p .

By Lemma 3.2, for each $n \geq 1$ we have $E[F^n] \cong \mu_{p^n}$ as I -modules. If we fix a torsion point $U \in E[F^k] \setminus E[F^{k-1}]$, then it follows that $T'' := \tau U - U$ has order p , so that $T' = rT''$ for some integer $0 < r < p$. Define $T := -rU$. Then

$$\tau(P + T) - (P + T) = (\tau P - P) - r(\tau U - U) = T' - rT'' = 0,$$

as desired. ■

Remark 3.7. The second part of the lemma can be interpreted in terms of Galois cohomology. Let G' be the cyclic group $\text{Gal}(\mathbb{Q}_p(\zeta_m)/\mathbb{Q}_p(\zeta_{m/p}))$. Since $E[F^k] \cong \mu_{p^k}$, the fact that for every $T' \in E[F]$ there exists $T \in E[F^k]$ such that $T' = \tau T - T$ is equivalent to the fact that the Galois cohomology group $H^1(G', \mu_m)$ is killed by m/p .

4. PROOF OF THEOREM 1.1

It is clear that in the proof of Theorem 1.1 we may replace K by a finite extension, so we assume from now on that K contains the field k of complex multiplication. Also, without loss of generality we may assume that E has everywhere good reduction over K (see [19, Theorem II.6.1]). This implies (see [19, Remark VI.4.1.1]) that for every finite extension L/K and every non-archimedean place w of L , we have $\lambda_w(Q) \geq 0$ for all nonzero points $Q \in E(L)$.

Let $D = [K : \mathbb{Q}]$. We choose once and for all (using the Chebotarev density theorem) a non-archimedean place v_0 of K of residue characteristic p satisfying the following conditions:

1. E has good reduction at v_0
2. p splits completely in K
3. $p > \exp(DC_1)$, where C_1 is the constant appearing in the statement of Corollary 2.3. (Equivalently, this inequality says that $A_p := (\log p)/D - C_1 > 0$.)

Note that since $k \subseteq K$, hypothesis (2) automatically implies that E has ordinary reduction at v_0 .

We will prove the following more explicit version of Theorem 1.1:

Theorem 4.1. *With K and p as above, let E/K be an elliptic curve having complex multiplication. Let C_1 be the constant from Corollary 2.3. Let L/K be an abelian extension, and let $P \in E(L)$ be a non-torsion point. If we set $A_p = (\log p)/D - C_1 > 0$, then*

$$\hat{h}(P) \geq \begin{cases} \frac{A_p}{2(p+1)} & \text{if } L/K \text{ is unramified above } v_0 \\ \frac{A_p}{4p} & \text{if } L/K \text{ is ramified above } v_0. \end{cases}$$

PROOF. We may assume without loss of generality that the extension L/K is finite. The proof proceeds by induction on the power of p dividing the local conductor at v_0 of L/K .

More concretely, fix a place w of L lying over v_0 , and let m be the smallest positive integer such that $L_w \subseteq \mathbb{Q}_p(\zeta_m)$. Such an m exists by the local Kronecker–Weber theorem, since $K_{v_0} \cong \mathbb{Q}_p$. If we write $m = p^r m'$, with $(p, m') = 1$, then we define the *local conductor* $\mathfrak{f}(L) = \mathfrak{f}_{v_0}(L/K)$ to be p^r . This coincides with the definition of the conductor from local class field theory. As the notation suggests, $\mathfrak{f}_{v_0}(L/K)$ is independent of the chosen place $w \mid v_0$.

Let $P \in E(L)$ be a non-torsion point. If $\mathfrak{f}(L) = 1$ (i.e., L/K is unramified above v_0), then by Lemma 3.3 (letting σ be the Frobenius element of L/K above v_0) we have

$$l_w(\sigma P - FP) \geq \log p$$

for all places w of L with $w|v_0$.

It follows that

$$\sum_{w|p} d_w l_w(\sigma P - FP) \geq \sum_{w|v_0} d_w \log p = d_{v_0} \log p = \frac{\log p}{D}.$$

As noted above, for every nonarchimedean place $w \in M_L^0$ and every nonzero point $Q \in E(L)$ we have

$$d_w l_w(Q) \geq 0.$$

By Corollary 2.3, we also know that for every archimedean place $w \in M_L^\infty$ we have

$$l_w(Q) \geq -C_1.$$

Applying these inequalities to $Q = \sigma P - FP$, and using the decomposition

$$\hat{h}(Q) = \sum_{v \in M_{\mathbb{Q}}} \sum_{w|v} d_w l_w(Q)$$

of the global canonical height as a sum of Néron local heights, we find that

$$\hat{h}(\sigma P - FP) \geq \frac{\log p}{D} - C_1 = A_p > 0.$$

Applying the parallelogram law

$$\hat{h}(P_1 - P_2) = 2\hat{h}(P_1) + 2\hat{h}(P_2) - \hat{h}(P_1 + P_2)$$

and noting that $\hat{h}(P_1 + P_2) \geq 0$, $\hat{h}(\sigma P) = \hat{h}(P)$, and $\hat{h}(FP) = p\hat{h}(P)$, we conclude that

$$\hat{h}(P) \geq \frac{A_p}{2(p+1)}.$$

This establishes the base case $\mathfrak{f}(L) = 1$ of the induction.

Assume now that $\mathfrak{f}(L) \geq p$ (so that L/K is ramified above v_0), and that the theorem is true for all abelian extensions L' of K such that $\mathfrak{f}(L') < \mathfrak{f}(L)$.

As the local conductor of $L(\zeta_m)$ over K is still $\mathfrak{f} = p^k$, we may assume without loss of generality that L contains the m th roots of unity, so that for all places w of L lying over v_0 we have $L_w \cong \mathbb{Q}_p(\zeta_m)$.

By Lemma 3.2, all points in $E[F^k]$ are defined over $\mathbb{Q}_p(\zeta_{p^k}) \subseteq \mathbb{Q}_p(\zeta_m) = L_w$. They are also defined over an abelian extension of K . Without loss of generality, we may therefore also assume that $K(E[F^k]) \subseteq L$.

Let $\tau \in \text{Gal}(L/K)$ be as in Lemma 3.6, and let $P' = \tau FP - FP$. If $P' \neq 0$, then by part (1) of Lemma 3.6 we have

$$l_w(P') \geq \log p$$

for all places w of L with $w|v_0$.

An argument parallel to that in the case $\mathfrak{f}(L) = 1$ now shows that

$$\hat{h}(P) \geq \frac{A_p}{4p}.$$

Finally, if $P' = 0$, then part (2) of Lemma 3.6 tells us that there exists a torsion point $T \in E[F^k] \subseteq E(L)$ such that $P + T$ is defined over the fixed field L' of τ . Note that $L'_w = \mathbb{Q}_p(\zeta_{m/p})$, and therefore $\mathfrak{f}(L') < \mathfrak{f}(L)$. Since $\hat{h}(P) = \hat{h}(P + T)$, we can apply the inductive hypothesis to $P + T$ to conclude. $\blacksquare$

5. PROOF OF THEOREM 1.4

Suppose E/K is an elliptic curve whose j -invariant is non-integral at the finite place v_0 of K . Without loss of generality, we may replace K by a finite extension and we henceforth assume that E/K is semistable, with split multiplicative reduction at v_0 . Let $\nu_0 = \text{ord}_{v_0}(j_E^{-1})$.

Suppose, for the sake of contradiction, that $\{P_i\}$ is an infinite sequence of points in $E(K^{\text{ab}})$ such that $\hat{h}(P_i) \rightarrow 0$. Let L_i be the Galois closure of $K(P_i)/K$, which by assumption is an abelian extension of K . Let e_i be the ramification index above v_0 of L_i/K .

We consider two cases.

Case 1: The sequence e_i is bounded above.

For notational convenience in the sequel, let e be an upper bound for the integers e_i , and define $c := \frac{1}{e^2\nu_0^2} > 0$. Pick any point Q from the sequence $\{P_i\}$, let L be the Galois closure of $K(Q)/K$, and let $Q_1, \dots, Q_N$ be the Galois conjugates of Q over K . Note that Northcott's theorem implies that we can choose Q so that N is as large as we please.

By repeatedly applying the parallelogram law

$$\hat{h}(Q_i + Q_j) + \hat{h}(Q_i - Q_j) = 2\hat{h}(Q_i) + 2\hat{h}(Q_j)$$

and noting that $\hat{h}(Q_i) = \hat{h}(Q_j) = \hat{h}(Q)$ and $\hat{h}(Q_i + Q_j) \geq 0$ for all i, j , we obtain (following [6]) the inequality

$$(1) \quad \hat{h}(Q) \geq \frac{1}{4N(N-1)} \sum_{i \neq j} \hat{h}(Q_i - Q_j).$$

In order to make use of this observation, we need to estimate the contribution at places lying over v_0 to the local decomposition of the global canonical height as a sum of Néron local heights.

To do this, let w be any place of L lying over v_0 . Then $\log |j_E|_w > 0$ and $\nu := \text{ord}_w(j_E^{-1}) = e(w/v_0)\nu_0 \leq e\nu_0$. By Lemma 2.4, we have, for each integer $N \geq 2$, the inequalities

$$(2) \quad \begin{aligned} \sum_{i \neq j} \lambda_w(Q_i - Q_j) &\geq \frac{1}{12} \log |j_E|_w \left(\left(\frac{N}{\nu} \right)^2 - N \right) \\ &\geq \frac{c}{12} N^2 \log |j_E|_w - \frac{N}{12} \log |j_E|_w, \end{aligned}$$

and therefore

$$(3) \quad \frac{1}{N(N-1)} \sum_{i \neq j} \lambda_w(Q_i - Q_j) \geq \frac{c}{12} \log |j_E|_w - \frac{1}{12(N-1)} \log |j_E|_w.$$

We have been working at the specific place v_0 , but by the same reasoning, we see that for any place w of L where E has bad (and therefore split multiplicative)

reduction we have

$$(4) \quad \frac{1}{N(N-1)} \sum_{i \neq j} l_w(Q_i - Q_j) \geq -\frac{1}{12(N-1)} \log |j_E|_w.$$

For any place w where E has good reduction, we have

$$(5) \quad \frac{1}{N(N-1)} \sum_{i \neq j} l_w(Q_i - Q_j) \geq 0.$$

Finally, for any infinite place w of L we have, by Lemma 2.2,

$$(6) \quad \frac{1}{N(N-1)} \sum_{i \neq j} l_w(Q_i - Q_j) \geq -\frac{1}{12(N-1)} \log^+ |j_E|_w + o(1),$$

where $o(1)$ denotes a term which tends to 0 as N tends to infinity.

Combining inequalities (2) through (6), and using the decomposition of the global height as a sum of local heights, we obtain, using Lemma 2.1:

$$(7) \quad \begin{aligned} \frac{1}{N(N-1)} \sum_{i \neq j} \hat{h}(Q_i - Q_j) &= \sum_{w \in M_L} d_w \frac{1}{N(N-1)} \left(\sum_{i \neq j} l_w(Q_i - Q_j) \right) \\ &\geq \frac{c}{12} \sum_{w|v_0} d_w \log |j_E|_w - \frac{1}{12(N-1)} \sum_{w \in M_L} d_w \log^+ |j_E|_w \\ &= \frac{c}{12} d_{v_0} \log |j_E|_{v_0} - \frac{1}{12(N-1)} h(j_E) \\ &= \frac{c}{12} d_{v_0} \log |j_E|_{v_0} + o(1). \end{aligned}$$

Combining this with inequality (1), we find that there are positive real constants $c' = d_{v_0} \log |j_E|_{v_0}/48$ and $c'' = cc'$ depending only on E, K, v_0 , and e such that

$$(8) \quad \hat{h}(Q) \geq c'' + o(1).$$

Since $d_i \rightarrow \infty$, Q can be chosen so that N is as large as we like. However, for N sufficiently large, the inequality (8) contradicts the fact that $h(P_i) \rightarrow 0$. We therefore conclude that Case 1 cannot occur.

Case 2: $e_i \rightarrow \infty$.

As before, we pick an element $Q \in E(L)$ from the sequence $\{P_i\}$. Let L be the Galois closure of $K(Q)/K$; since L/K is abelian, the inertia groups of L/K at each place lying over v_0 are all the same group I .

Let $Q_1, \dots, Q_N$ be the conjugates of Q under I . Note that the hypothesis $e_i \rightarrow \infty$ guarantees that we can choose Q so that N is as large as we like.

Since the Q_i 's are conjugate under the inertia group I , we see that for every place w of L dividing v_0 , $Q_i - Q_j$ is in the kernel of reduction mod w for all i, j . It follows that for all $i \neq j$ we have

$$\lambda_w(Q_i - Q_j) \geq \frac{1}{12} \log |j_E|_w$$

and therefore

$$\frac{1}{N(N-1)} \sum_{i \neq j} \lambda_w(Q_i - Q_j) \geq \frac{1}{12} \log |j_E|_w$$

for all $w \mid v_0$.

Using the parallelogram law and Lemma 2.1 as before, we find that

$$\begin{aligned}\hat{h}(Q) &\geq \frac{1}{4} \sum_{w|v_0} d_w \left(\frac{1}{12} \log |j_E|_w \right) + o(1) \\ &= \frac{1}{48} d_{v_0} \log |j_E|_{v_0} + o(1) \\ &= c' + o(1),\end{aligned}$$

which for N large contradicts the fact that $\hat{h}(P_i) \rightarrow 0$. ■

ACKNOWLEDGEMENTS

The author would like to thank Brian Conrad for helpful discussions and Robert Rumely for his useful comments on an earlier version of this manuscript.

REFERENCES

- [1] F. Amoroso and R. Dvornicich, *A lower bound for the height in abelian extensions*, J. Number Theory **80** (2000), no. 2, 260–272.
- [2] F. Amoroso, U. Zannier, *A relative Dobrowolski lower bound over abelian extensions*. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **29** (2000), no. 3, 711–727.
- [3] M. Artin, Néron models, in *Arithmetic Geometry*, ed. G. Cornell and J. Silverman, Springer–Verlag, Berlin and New York, 1986.
- [4] E. Bombieri and U. Zannier, *A note on heights in certain infinite extensions of $\mathbf{Q}$* , Rend. Mat. Acad. Lincei, s. 9 **12** (2001), 5–14.
- [5] G. Faltings, *Complements to Mordell*, in G. Faltings, G. Wüstholz et. al, *Rational points*, Vieweg, 1984.
- [6] M. Hindry and J. Silverman, *On Lehmer’s conjecture for elliptic curves*, in Séminaire de Théorie des Nombres, Paris (1988–1989), 103–116, Progr. Math. 91, Birkhuser Boston, Boston, MA, 1990.
- [7] M. Hindry and J. Silverman, *Sur le nombre de points de torsion rationnels sur une courbe elliptique*, C. R. Acad. Sci. Paris Sér. I Math. **329** (1999), no. 2, 97–100.
- [8] S. Lang, *Arakelov Theory*, Springer–Verlag, Berlin and New York, 1988.
- [9] S. Lang, *Elliptic Functions*, Springer–Verlag, Berlin and New York, 1987.
- [10] T. Oda, *The first de Rham cohomology group and Dieudonné modules* Ann. Sci. École Norm. Sup. (4) **2** (1969), 63–135.
- [11] B. Poonen, *Mordell–Lang plus Bogomolov*, Invent. Math. **137** (1999), 413–425.
- [12] K. Ribet, appendix to *Finiteness theorems in geometric classfield theory* by N. Katz and S. Lang, Enseign. Math. (2) **27** (1981), no. 3–4, 285–319.
- [13] W. Ruppert, *Torsion points of abelian varieties in abelian extensions*, preprint. (1998).
- [14] J–P. Serre, *Propriétés galoisiennes des points d’ordre fini des courbes elliptiques*, Invent. Math. **15** (1972), no. 4, 259–331.
- [15] J–P. Serre, *Lectures on the Mordell–Weil Theorem*, Aspects of Mathematics, Vieweg and Sohn, Braunschweig, 1997.
- [16] J–P. Serre and J. Tate, *Good reduction of abelian varieties*, Ann. of Math. (2) **88** (1968), 492–517.
- [17] J. Silverman, *Lower bound for the canonical height on elliptic curves*, Duke Math. J. **48** (1981), 633–648.
- [18] J. Silverman, *The Arithmetic of Elliptic Curves*, Springer–Verlag, Berlin and New York, 1986.
- [19] J. Silverman, *Advanced Topics in the Arithmetic of Elliptic Curves*, Springer–Verlag, Berlin and New York, 1994.
- [20] L. Szpiro, E. Ullmo, S. Zhang, *Équirépartition des petits points*, Invent. Math. **127** (1997), no. 2, 337–347.
- [21] J. Tate, Finite Flat Group Schemes, in *Modular Forms and Fermat’s Last Theorem*, ed. G. Cornell, J. Silverman, and G. Stevens, Springer–Verlag, Berlin and New York, 1997.
- [22] L. Washington, *Cyclotomic Fields* (2nd edition), Springer–Verlag, Berlin and New York, 1997.
- [23] J. Zarhin, *Endomorphisms and torsion of abelian varieties*, Duke Math. J. **54** (1987), 131–145.
- [24] S. Zhang, *Equidistribution of small points on abelian varieties*, Ann. of Math. (2) **147** (1998), no. 1, 159–165.

E-mail address: `mbaker@math.uga.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GEORGIA, ATHENS, GA 30602-7403